



Berlot, B., Harries, I., & Bucciarelli-Ducci, C. (2019). Connection between the heart and the gut. *Heart*. <https://doi.org/10.1136/heartjnl-2019-314832>

Peer reviewed version

License (if available):  
CC BY-NC

Link to published version (if available):  
[10.1136/heartjnl-2019-314832](https://doi.org/10.1136/heartjnl-2019-314832)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via BMJ Publishing at <https://heart.bmj.com/content/early/2019/04/08/heartjnl-2019-314832> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

## The connection between the heart and the gut

B. Berlot<sup>1,2</sup>, I. Harries<sup>1,2</sup>, C. Bucciarelli – Ducci<sup>1,2,3</sup>

<sup>1</sup>Bristol Heart Institute, University Hospitals Bristol

<sup>2</sup>Bristol Medical school, University of Bristol

<sup>3</sup>Bristol NIHR Biomedical Research Centre, University Hospitals Bristol NHS Trust and University of Bristol. Bristol, United Kingdom.

**CLINICAL PROBLEM:** A 45-year old male with ulcerative colitis was admitted with bloody diarrhoea and chest pain. Inflammatory markers and high-sensitivity troponin were elevated (CRP 57 mg/L, WBC  $10.65 \times 10^9/L$ , Neutrophil  $6.6 \times 10^9/L$ , TnI 663 mmol/L). The electrocardiogram (ECG) showed inferior ST- elevation. Urgent coronary angiography revealed unobstructed coronary arteries. Inpatient cardiovascular magnetic resonance (CMR) was arranged to determine the aetiology of the myocardial infarction with non-obstructive coronary arteries (MINOCA). The imaging protocol at 1.5 T included balanced steady state free precession (bSSFP) cine images, T2-weighted oedema sequences, early and late gadolinium enhancement (LGE). Native T1 and T2 mapping images provided advanced tissue characterisation.

**QUESTION:** What is the most likely diagnosis based on the MRI findings?

- A) Multiple embolic myocardial infarctions in the right coronary artery territory
- B) Acute autoimmune myocarditis
- C) Cardiac sarcoidosis
- D) Stress (Takotsubo) cardiomyopathy
- E) Multiple embolic myocardial infarctions in the left circumflex coronary artery territory

**ANSWER:** B) Multiple patchy, mid-myocardial, high-intensity signal areas in the inferolateral wall on T2 weighted images (B) and increased T2 mapping values on the short axis images (G) are highly specific for oedema. These areas have high-signal intensity on early (C) and late (D,H) sequences, suggesting hyperaemia with capillary leakage and increased extracellular space, respectively. Furthermore, increased native T1 mapping values in all layers but predominantly in the subepicardium of the entire lateral wall (F) are in keeping with extensive myocardial injury. Taken in combination, these findings meet the updated Lake Louise criteria for acute myocarditis (1). Despite extensive oedema, stress (Takotsubo) cardiomyopathy (Option D) can be excluded by ventricular morphology and presence of LGE (2), and multiple embolic myocardial infarctions (Option A and E) are excluded by the non-ischemic pattern of LGE (3). Finally, based on the CMR, cardiac sarcoidosis (Option C) cannot be easily distinguished from acute autoimmune myocarditis. However, the clinical presentation and absence of pulmonary changes make the latter diagnosis the most likely (4). Cardiac biopsy might definitively differentiate, but is not routinely performed due to associated procedural risk. The patient was treated with corticosteroids and monoclonal antibodies and responded well. The pathophysiological mechanisms underlying cardiac manifestations of inflammatory bowel disease are incompletely understood but may occur in up to one third of patients with inflammatory bowel disease (5).

1. Ferreira VM et al. Cardiovascular Magnetic Resonance in Nonischemic Myocardial Inflammation: Expert Recommendations. *J Am Coll Cardiol* 2018Dec;72(24):3158–76.
2. Plácido R, et al. The role of cardiovascular magnetic resonance in takotsubo syndrome. *J Cardiovasc Magn Reson* 2016 Oct;18(1):68.
3. Mahrholdt H, et al. Delayed enhancement cardiovascular magnetic resonance assessment of non-ischaemic cardiomyopathies. *Eur Heart J. England*; 2005 Aug;26(15):1461–74.
4. Kadkhodayan A, et al. Imaging of Inflammation in Unexplained Cardiomyopathy. *JACC Cardiovasc Imaging* 2016 May;9(5):603–17.
5. Greenstein, et al. The extraintestinal complications of Crohn's disease and ulcerative colitis: A study of 700 patients. *Medicine (Baltimore)* 1976; 55:401-412.

### **Acknowledgment**

CBD is in part supported by the NIHR Biomedical Research Centre at University Hospitals Bristol NHS Foundation Trust and the University of Bristol. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health and Social Care.

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non-exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in HEART editions and any other BMJ PGL products to exploit all subsidiary rights

